


CENTRAL QUEENSLAND SEISMOLOGY RESEARCH GROUP (CQSRG)




# CQSRG Seismological Report 2018

Edition 1.01

Michael Turnbull BAppSc(Distinction) MAppSc,  
*Lead Seismologist, CQSRG.*

Version 1



RESEARCHING EARTHQUAKES IN CENTRAL QUEENSLAND SINCE 2002.

Web site: <http://cqsr.org>

## Table of Contents

Edition control.....	iii
Version Changes.....	iii
Versions 1.00.....	iii
Copyright.....	iv
Disclaimer.....	iv
Introduction .....	1
CQSRG Station Reports .....	2
FS03 Uptime 2018.....	2
WOOW Uptime 2018 .....	2
The Continuing Bowen Earthquake Sequence.....	3
CQSRG Main Earthquake Catalogue 2018 .....	10
2018 Statistical Summary.....	16
2018 Earthquake Maps.....	18
Public Seismic Network (PSN) .....	20
Appendix A – Details of FS03 .....	21
Appendix B – Details of WOOW.....	22
Appendix C – CQSRG Method of Earthquake Location.....	23
Appendix D – CQSRG Method of Magnitude Quantification from FS03 Records.....	24
Calibration of FS03 Seismometer for Earthquake Magnitude Determination. ....	24
Introduction .....	24
Background Information .....	24
Method .....	25
Important Note Concerning Accuracy and Precision.....	26
Example Usage.....	27
Student Resources .....	28
Appendix E - CQSRG Method of Magnitude Quantification from EIDS Records .....	31
Relative Calibration of EIDS Seismometer for Earthquake Magnitude Determination Based on FS03 Past events.....	31
Introduction .....	31
Background Information .....	31
Method .....	31
Example Usage.....	34

Appendix F - Magnitude Calibration of BW1H for the Bowen 2016 Earthquake Sequence.....	35
Introduction .....	35
Method of Calibration.....	36
Verification of Calibration .....	37
Calibration Saturation .....	37
Appendix G - Method Used to Identify Bowen August 2016 ML 5.8 Aftershocks.....	38
Analysis of S-P Times.....	38
The characteristic shape of the wave form .....	39

## **Edition control**

Date of Release	Edition Number	Comments
March 2021	1.00	The original edition.
March 2021	1.01	Corrected internal Figure link errors and incorrect year error.

## **Version Changes**

### **Version 1.00**

The original version.

## Copyright

©Copyright CQSRG 2021. All rights reserved. A licence is issued to institutes to freely use and copy this document for non-commercial purposes in its unaltered state.

This document may be cited as follows.

(CQSRG 2021), *CQSRG Seismological Report 2018* Central Queensland Seismology Research Group, Author: Michael Turnbull, Published March 2021 by Email distribution and online at <http://cqsrg.org/reports/>

## Disclaimer

Every effort has been made to ensure that the information contained in this document is accurate, in the author's opinion, at the time the document was created; however, the author is human, and therefore subject to human frailty, and may well have made several errors in judgement in this document. Therefore the author cannot guarantee that the information contained in this document is absolutely accurate. The information contained in this document consists of the scientific opinions of the author, and may not coincide with the opinions of the reader. Therefore, the information contained in this document is advisory only, and not absolute. Basically this means that if you agree with the opinions of the author then that is your decision, as it is if you disagree with the author's opinions.

Mike Turnbull may be contacted by email at: [Mike.Turnbull@cqsrg.org](mailto:Mike.Turnbull@cqsrg.org)

## Introduction

This report details earthquakes detected and located by the Central Queensland Seismology Research Group (CQSRG) during the 2018 calendar year. Technical and statistical summaries of earthquakes that occurred in Queensland are provided. The date and time of earthquakes noted in this report are provided in Universal Coordinated Time (UTC).

Data and information provided in this report may supersede or supplement data and information provided in previous CQSRG Annual Seismological Reports. This is due to ongoing CQSRG research that may add to or revise data and information collected and analysed from previous years.

CQSRG was established in 2002, under the auspices of the Faculty of Informatics and Communication of Central Queensland University (CQU), with Michael Turnbull (Lecturer, and later Adjunct Research Fellow) and Kevin McCue (Visiting Professor, and later Adjunct Professor) as the designated researchers. This affiliation with CQU continued until February 2013, when, due to a divergence in academic focus of CQU and CQSRG, the researchers allowed their Adjunct appointments to lapse. From February 2013 until December 2016, CQSRG operated independently of CQU, with the same two people conducting the research. In mid-2016 Dr Andrew Hammond, Senior Lecturer in Geology at CQU, joined CQSRG as a research collaborator. Mike Turnbull's and Kevin McCue's adjunct academic appointments with CQU were re-established in October 2016.

During the 2018 calendar year CQSRG operated two seismic monitoring stations, designated FS03, and WOOW. Details of these stations, including location and equipment, are provided in Appendix A. This report contains information relating to earthquakes detected by the FS03 and WOOW seismic monitoring stations, as well as earthquakes of significance located within Queensland, but outside the CQSRG detection area, and reported by Geoscience Australia (GA).

The WOOW monitoring station was decommissioned on 2018-03-06 18:00 (UTC).

CQSRG locates and quantifies earthquakes using the methods detailed in Appendices D, E and F.

## CQSRG Station Reports

### FS03 Uptime 2018




Figure 1: Percentage Uptime/Downtime of FS03 during 2018

The FS03 station has been in continuous operation since 2003-01-01 00:00:00.00 (UTC). Technical details of the station are found at <http://cqsrg.org/network/FS03technical/>.

Throughout the 2018 calendar year the FS03 station was actively monitoring for seismic events greater than 98% of the year.

This high proportion of availability was due to automation of the data download process, and the provision of an RS232 serial data radio link to the station in April 2015.

The down time was entirely comprised of data download time.




Figure 2: Percentage Uptime/Downtime of WOOW during 2017.

### WOOW Uptime 2018

The WOOW station was installed in December 2016, and commissioned on 2017-01-10 03:10 (UTC). Technical details of the station are found at <http://cqsrg.org/network/WOOWtechnical/>. It was decommissioned on 2018-03-06 18:00 (UTC).

In the 2018 calendar year, up to the time it was decommissioned, the WOOW station was actively monitoring for seismic events less than 4% of the time.

This high proportion of unavailability was due to failures of the Telstra mobile data modem link from January to March combined with an inability to service the station.

During 2018 the records obtained from the WOOW station made no contribution to the detection and location of earthquakes in the Central Queensland region.

The station will not be re-established in the future by CQSRG. The housing and concrete pad have been left in place, and it has 240V mains power available in the housing.

## The Continuing Bowen Earthquake Sequence

On 2016-08-18 at 04:30 UTC an ML 5.8 earthquake<sup>1</sup> occurred 63 km north east of Bowen in the Whitsunday Passage. This was followed over the next three weeks by 77 aftershocks ranging from ML 1.6 to ML 4.2 that were located by CQSRG; however there were many more aftershocks of magnitudes below ML 1.6 that have been identified by CQSRG in the Bowen Urban Monitoring (UM) network site (BW1H) seismic records but have not been located due to insufficient recordings. This sequence of events is known within CQSRG as the **2016 Bowen Earthquake** and Aftershock Sequence (BW 2016).




Figure 3: Bowen aftershocks that occurred in 2018, and were located by CQSRG, in relation to the August 2016 ML 5.8 earthquake (shown as orange marker).

During 2018 calendar year 189 aftershocks ranging in magnitude from ML 2.8 down to ML 0.7 were detected by CQSRG; not all of which were located, due to lack of sufficient instrumental records.

Of those events that were detected, 30 provided sufficient recordings to be reliably located. A map showing locations of aftershocks that were located by CQSRG during 2018 is presented in Figure 3. This figure also shows the location of the main ML 5.8 August 2016 event.

Figure 4 shows a graphical plot of the 2016 Bowen sequence up to the end of 2018.

<sup>1</sup> The magnitude assigned by CQSRG is that assigned by Geoscience Australia.






Figure 4: Events associated with the 2016 Bowen ML 5.8 earthquake, as detected and located by CQSRG.

It is clear from Figure 4 that the Bowen 2016 sequence is ongoing; and all indications are that it will continue into the future.

CQSRG has also inspected Bowen regional seismic records going back to the beginning of 2015 to detect any earthquake events that occurred in the area prior to the main August 2016 earthquake; and these are also shown in Figure 4. Although there were a few (six) events detected and located (ranging in magnitude from ML 1.4 to ML 3.6) it is not clear whether these were precursor events, or if they are simply representative of the pre-August 2016 background seismicity.

Extending the investigation further back in time will prove difficult due to lack of reliable seismic records.

A full listing of the 189 Bowen earthquake events detected by CQSRG in 2018 is given in Table 1. Due to lack of available data not all of the listed events had mathematical locational analysis performed.

The method used to identify Bowen August 2016 ML 5.8 aftershocks is detailed in Appendix G - Method Used to Identify Bowen August 2016 ML 5.8 Aftershocks

Table 1: Earthquake events in the Whitsunday Passage area detected by CQSRG in 2018.

Magnitude (ML)	Date (UTC)	Time (UTC)	Latitude	Longitude
2.8	2018-02-03	03:43:10.48	-19.870	148.770
2.7	2018-06-07	12:05:59.80	-20.313	148.747
2.6	2018-02-10	16:47:04.75	-20.105	148.780

<b>Magnitude (ML)</b>	<b>Date (UTC)</b>	<b>Time (UTC)</b>	<b>Latitude</b>	<b>Longitude</b>
2.3	2018-01-09	10:24:25.89	-20.160	148.775
2.2	2018-01-25	17:37:02.44	-19.742	148.739
2.2	2018-02-18	19:21:55.83	-19.917	148.775
2.2	2018-04-10	08:57:20.63	-19.610	148.721
2.2	2018-05-14	01:07:04.08	-19.731	148.493
2.2	2018-07-28	10:54:40.99	-20.115	148.829
2.2	2018-08-08	01:26:22.10	-20.199	148.801
2.2	2018-10-22	20:27:04.62	-19.800	148.779
2.1	2018-04-23	04:43:54.20	-19.989	148.836
2.1	2018-08-13	10:11:44.92	-20.054	148.810
2.1	2018-09-14	23:23:39.37	-19.743	148.522
2.1	2018-09-21	22:11:55.69	-20.167	148.778
2.1	2018-10-02	20:15:59.37	-20.089	148.886
2.1	2018-11-27	20:47:49.35	-20.106	148.811
2.0	2018-01-08	14:43:03.03	-20.248	148.715
2.0	2018-02-04	18:37:01.81	-20.126	148.801
2.0	2018-04-05	09:51:24.38	-19.992	148.763
2.0	2018-04-05	11:27:49.73	-20.063	148.841
2.0	2018-05-08	10:54:34.51	-20.030	148.946
2.0	2018-05-16	17:45:58.24	-20.044	148.784
2.0	2018-05-25	13:43:29.75	-20.016	148.755
2.0	2018-06-07	23:01:37.68	-19.898	148.775
2.0	2018-08-11	05:07:03.19	-20.103	148.787
2.0	2018-12-22	21:15:13.98	-19.830	148.774
1.9	2018-04-23	21:56		
1.9	2018-04-23	03:46		
1.9	2018-01-11	12:28		
1.9	2018-07-11	06:19:52.05	-20.106	148.691
1.9	2018-12-22	22:32		
1.9	2018-12-23	21:15		
1.9	2018-12-25	12:27		
1.9	2018-01-23	07:22		
1.9	2018-08-08	10:45		
1.8	2018-08-22	21:48		
1.8	2018-02-20	15:27		
1.8	2018-07-23	03:56		
1.8	2018-03-30	18:31		
1.8	2018-08-02	10:20		
1.8	2018-09-16	08:54		
1.8	2018-03-04	11:59		
1.8	2018-02-16	22:25:13.45	-19.971	148.784
1.8	2018-11-19	05:37		
1.8	2018-07-31	16:00		

<b>Magnitude (ML)</b>	<b>Date (UTC)</b>	<b>Time (UTC)</b>	<b>Latitude</b>	<b>Longitude</b>
1.8	2018-01-04	18:37		
1.8	2018-09-29	02:32		
1.8	2018-08-11	01:31		
1.8	2018-01-01	09:05		
1.8	2018-08-13	10:02		
1.8	2018-03-28	21:00		
1.8	2018-08-21	16:16		
1.8	2018-10-03	19:04		
1.7	2018-01-20	02:20		
1.7	2018-09-25	17:26		
1.7	2018-04-18	14:07		
1.7	2018-08-09	08:30		
1.7	2018-10-03	19:07		
1.7	2018-01-31	09:48		
1.7	2018-02-18	00:57		
1.7	2018-08-11	06:22		
1.7	2018-07-28	03:57		
1.7	2018-10-14	14:57		
1.7	2018-01-09	15:07:37.09	-20.074	148.831
1.7	2018-10-30	15:25		
1.7	2018-11-30	11:32		
1.7	2018-12-02	06:29		
1.7	2018-12-29	18:03		
1.7	2018-04-20	08:07		
1.7	2018-05-09	21:07		
1.7	2018-10-09	13:38		
1.7	2018-01-19	11:21		
1.7	2018-05-11	13:35		
1.7	2018-03-21	11:33		
1.7	2018-04-16	14:42		
1.7	2018-05-09	14:19		
1.7	2018-03-26	19:32		
1.7	2018-01-20	11:52		
1.7	2018-04-16	08:23		
1.7	2018-03-01	23:58		
1.7	2018-06-02	05:28		
1.7	2018-01-31	11:04		
1.6	2018-02-11	13:29		
1.6	2018-08-21	21:00		
1.6	2018-03-15	07:25		
1.6	2018-03-27	07:37		
1.6	2018-08-11	09:09		
1.6	2018-03-19	19:16		

<b>Magnitude (ML)</b>	<b>Date (UTC)</b>	<b>Time (UTC)</b>	<b>Latitude</b>	<b>Longitude</b>
1.6	2018-01-28	17:45		
1.6	2018-05-01	09:49		
1.6	2018-07-31	20:17		
1.6	2018-05-31	06:05		
1.6	2018-04-22	03:40		
1.6	2018-01-25	10:06		
1.6	2018-04-08	14:43		
1.6	2018-06-14	22:26		
1.6	2018-02-12	18:19		
1.6	2018-05-22	11:12		
1.6	2018-08-13	16:53		
1.6	2018-02-02	09:00		
1.6	2018-10-31	06:18		
1.6	2018-11-09	17:28		
1.6	2018-11-13	21:25		
1.6	2018-11-18	18:07		
1.6	2018-12-01	00:00		
1.6	2018-12-17	23:14		
1.6	2018-06-15	02:49		
1.6	2018-02-27	15:50		
1.6	2018-01-08	12:29		
1.6	2018-04-15	09:23		
1.6	2018-03-01	16:46		
1.6	2018-04-05	18:27		
1.6	2018-01-06	06:23		
1.6	2018-04-15	15:16		
1.6	2018-08-31	22:32		
1.6	2018-07-30	21:54		
1.6	2018-03-24	16:22		
1.6	2018-08-11	20:27		
1.6	2018-01-26	19:24		
1.6	2018-08-06	19:52		
1.6	2018-02-20	15:11		
1.6	2018-08-10	18:58		
1.6	2018-01-25	12:14		
1.6	2018-02-23	08:38		
1.6	2018-01-31	14:38		
1.6	2018-08-17	13:18		
1.5	2018-02-02	20:34		
1.5	2018-01-21	16:18		
1.5	2018-06-11	19:25		
1.5	2018-03-25	04:24		
1.5	2018-08-29	23:17		

<b>Magnitude (ML)</b>	<b>Date (UTC)</b>	<b>Time (UTC)</b>	<b>Latitude</b>	<b>Longitude</b>
1.5	2018-03-05	08:09		
1.5	2018-05-10	16:33		
1.5	2018-10-03	13:10		
1.5	2018-04-15	02:54		
1.5	2018-08-10	17:00		
1.5	2018-02-10	11:53		
1.5	2018-05-02	17:38		
1.5	2018-10-30	10:49		
1.5	2018-11-26	09:50		
1.5	2018-12-01	21:03		
1.5	2018-12-03	19:11		
1.5	2018-12-10	03:57		
1.5	2018-12-22	21:35		
1.5	2018-06-01	23:37		
1.5	2018-01-28	10:18		
1.5	2018-01-28	09:09		
1.5	2018-08-25	19:37		
1.5	2018-04-24	07:29		
1.5	2018-09-17	01:49		
1.5	2018-03-24	19:41		
1.5	2018-04-30	12:11		
1.5	2018-01-06	14:34		
1.5	2018-01-28	10:34		
1.5	2018-06-13	23:18		
1.5	2018-05-31	10:27		
1.5	2018-10-09	13:38		
1.5	2018-02-10	19:29		
1.5	2018-04-24	17:32		
1.5	2018-10-21	09:56		
1.5	2018-08-06	09:34		
1.5	2018-06-09	13:18		
1.5	2018-01-26	17:23		
1.4	2018-06-02	19:42		
1.4	2018-10-09	13:38		
1.4	2018-03-19	17:08		
1.4	2018-03-12	08:38		
1.4	2018-08-17	17:01		
1.4	2018-07-31	13:51		
1.4	2018-01-22	05:17		
1.4	2018-08-29	11:00		
1.4	2018-03-01	18:15		
1.4	2018-11-07	10:40		
1.4	2018-11-11	20:02		

<b>Magnitude (ML)</b>	<b>Date (UTC)</b>	<b>Time (UTC)</b>	<b>Latitude</b>	<b>Longitude</b>
1.4	2018-11-15	15:26		
1.4	2018-11-26	12:06		
1.4	2018-11-29	16:28		
1.4	2018-12-18	12:45		
1.4	2018-12-22	09:10		
1.4	2018-01-01	09:40		
1.4	2018-06-11	10:04		
1.4	2018-08-30	17:51		
1.4	2018-09-27	12:45		
1.4	2018-08-19	13:25		
1.4	2018-03-27	12:31		
0.9	2018-12-03	08:47		
0.7	2018-10-31	14:20		
0.7	2018-11-17	18:45		

The available evidence indicates that the BW 2016 aftershock sequence is continuing, and will continue into the foreseeable future. What is currently being observed may well represent the initiation of a new long-term seismicity regime for the Whitsunday Passage area.

Although the unlocated events listed in Table 1 are not included in the Main CQSRG Earthquake Catalogue, CQSRG has the source data used to identify those events on record, and this data can be made available to interested parties on request.

## **CQSRG Main Earthquake Catalogue 2018**

During 2018, 75 earthquake events were detected, located, and catalogued by CQSRG. Details of these events are provided in Table 2. The online full version of the CQSRG catalogue can be accessed at <http://cqsrg.org/catalogue/>. The 75 events listed in Table 2 include the 30 Bowen aftershock events that were sufficiently well recorded to have been located.

An additional 159 Bowen aftershock events were identified, but insufficient recordings of those events were available to allow for reliable locations. While the methodology used to identify BW 2016 aftershocks is considered reliable, it is possible that a small number of them (less than 3%) have been incorrectly assigned. For this reason, these 159 events have not been included in the main CQSRG Earthquake Catalogue, but are included in a CQSRG Supplementary Catalogue, listed in Table 1, and shown in Figure 4.

It should also be noted that, although the main ML 5.8 Bowen event and some of the listed aftershocks were well recorded on the CQSRG FS03 station, numerous other aftershocks that are listed in the CQSRG main and supplementary catalogues were not principally detected on the CQSRG network. Most of those aftershock events were identified by manual inspection of the daily records obtained off the BW1H and BW2S Urban Monitoring (UM) stations at Bowen.

It is also noted that, where the EQLOCL algorithm could not calculate a depth due to lack of vertical resolution, the focal depths listed in the CQSRG Earthquake Catalogue have been constrained to the local norm (10 km).

Table 2: Earthquake Events Detected, Located, and Catalogued by CQSRG during 2018.

Date (UTC)	Time (UTC)	Longitude	Latitude	Depth (km)	Magnitude (ML)	Place	Comment
2018-12-25	19:33:58.91	151.722	-25.155	10	0.3	Mt Perry	SW FS03. 8 km NE Mt Perry. Reviewed 2019-01-23.
2018-12-22	21:15:13.98	148.774	-19.830	10	2.0	Bowen	Aftershock of 2016-08-18 04:30 event. 62 km NE Bowen. Reviewed 2019-01-24.
2018-11-29	06:10:20	151.725	-25.155	10	1.5	Mt Perry	9 km NE Mt Perry. Reviewed 2018-12-02.
2018-11-27	20:47:49.35	148.811	-20.106	10	2.1	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 20 km NE Airlie Beach. Reviewed 2019-01-24.
2018-11-24	23:54:10.61	151.473	-25.239	10	1.0	Mt Perry	18 km SW Mt Perry. Reviewed 2018-11-27.
2018-11-21	01:15:30	151.867	-25.107	10	1.0	FS03	Within 20 km of FS03. Not recorded on EIDS therefore probably to the NE quadrant of FS03 between Gin Gin and Wallaville. Reviewed 2018-11-27.
2018-11-17	20:52:03	151.867	-25.107	10	1.1	FS03	Within 10 km of FS03. Not recorded on EIDS therefore probably to the NE quadrant of FS03 between Gin Gin and Wallaville. Reviewed 2018-11-18.
2018-11-08	13:43:38.79	147.800	-20.018	10	1.8	Bowen	Related to the 2011-04-16 05:31 ML5.3 earthquake? 47 km W Bowen. Reviewed 2018-11-16.
2018-11-06	14:50:23.02	151.746	-25.147	10	0.7	Mt Perry	10 km NE Mt Perry. Reviewed 2018-11-16.
2018-11-03	21:46:06.03	151.513	-25.226	10	1.5	Mt Perry	14 km SW Mt Perry. Reviewed 2018-11-04.
2018-10-24	10:38:08.19	151.708	-24.832	10	1.1	Gin Gin	30 km NW Gin Gin. Reviewed 2018-11-04.
2018-10-22	20:27:04.62	148.779	-19.800	10	2.2	Bowen	Aftershock of 2016-08-18 04:30 event. 60 km NE Bowen. Reviewed 2019-01-24.
2018-10-02	20:15:59.37	148.886	-20.089	10	2.1	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 26 km NE Airlie Beach. Reviewed 2018-10-03.
2018-09-30	03:11:12.33	151.534	-25.219	10	1.5	Mt Perry	12 km SW Mt Perry. Reviewed 2018-10-03.
2018-09-27	23:22	151.867	-25.107	10	1.0	Gin Gin/ Wallaville	Gin Gin/Wallaville. Only recorded on FS03. First motion is from the NE. Reviewed 2018-10-03.
2018-09-26	13:09:19.67	151.853	-25.373	10	1.6	Biggenden	24 km NW Biggenden. Reviewed 2018-10-05.
2018-09-21	22:11:55.69	148.778	-20.167	10	2.1	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 13 km NE Airlie Beach. Reviewed 2018-09-22.



Date (UTC)	Time (UTC)	Longitude	Latitude	Depth (km)	Magnitude (ML)	Place	Comment
2018-09-20	21:27:39.98	151.783	-25.510	10	0.9	Ban Ban Springs	19 km N Ban Ban Springs. Viewed 2018-09-21.
2018-09-14	23:23:39.37	148.522	-19.743	10	2.1	Bowen	Aftershock of 2016-08-18 04:30 event. 41 km NE Bowen. Reviewed 2018-09-15.
2018-09-06	10:02:24.24	150.767	-25.577	10	2.0	Nerangy	42 km SW Eidsvold.8 km N Nerangy. Reviewed 20180906. Not heard or felt at Nerangy.
2018-08-13	10:11:44.92	148.810	-20.054	10	2.1	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 26 km NE Airlie Beach. Reviewed 2018-09-23.
2018-08-11	05:07:03.19	148.787	-20.103	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 20 km NE Airlie Beach. Reviewed 2018-09-23.
2018-08-08	23:09:56.85	151.492	-25.233	10	0.8	Mt Perry	16 km SW Mt Perry. Reviewed 2018-08-13.
2018-08-08	01:26:22.10	148.801	-20.199	10	2.2	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 12 km NE Airlie Beach Reviewed 2018-08-14.
2018-08-06	21:25:09.76	153.684	-23.153	10	2.5	Coral Sea	214 km NE Seventeen Seventy. Reviewed 2018-08-13.
2018-08-05	18:32:12.68	151.672	-25.172	10	0.8	Mt Perry	3 km NE Mt Perry. Reviewed 2018-08-13.
2018-07-28	10:54:40.99	148.829	-20.115	10	2.2	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 21 km NE Airlie Beach Reviewed 2018-07-30.
2018-07-17	15:28:39.25	151.645	-24.739	10	1.6	Gin Gin	42 km NW Gin Gin. Reviewed 2018-07-18.
2018-07-16	12:32:42.96	140.815	-25.700	10	3.1	Betoota	7 km E Betoota and Mt Leonard Station. Reviewed 2018-07-17.
2018-07-15	19:07:06.30	151.543	-25.216	10	0.7	Mt Perry	Aftershock of 2018-06-28 14:33 event. 11 km SW Mt Perry. Reviewed 2018-07-16.
2018-07-11	06:19:52.05	148.691	-20.106	10	1.9	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 18 km N Airlie Beach Reviewed 2018-07-16.
2018-06-28	14:33:42.61	151.541	-25.216	10	0.9	Mt Perry	11 km SW Mt Perry. Reviewed 2018-07-04.
2018-06-07	23:01:37.68	148.775	-19.898	10	2.0	Bowen	Aftershock of 2016-08-18 04:30 event. 57 km NE Bowen Reviewed 2018-06-08.
2018-06-07	12:05:59.80	148.747	-20.313	10	2.7	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 6 km SE Airlie Beach Reviewed 2018-06-08.

Date (UTC)	Time (UTC)	Longitude	Latitude	Depth (km)	Magnitude (ML)	Place	Comment
2018-05-28	01:36:22.34	154.591	-22.389	10	4.1	Coral Sea	404 km NE Gladstone, on the continental drop off.
2018-05-26	11:09:05.19	150.784	-24.703	10	2.0	Monto	39 km NW Monto. Reviewed 2018-05-29.
2018-05-25	13:43:29.75	148.755	-20.016	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 28 km N Airlie Beach Reviewed 2018-05-31.
2018-05-16	17:45:58.24	148.784	-20.044	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 25 km N Airlie Beach Reviewed 2018-05-31.
2018-05-14	01:07:04.08	148.493	-19.731	10	2.2	Bowen	Aftershock of 2016-08-18 04:30 event. 48 km NE Bowen. Reviewed 2018-04-15.
2018-05-10	00:52:15.69	150.981	-24.127	10	2.0	Calliope	26 km SW Calliope. Reviewed 2018-05-10. Not a blast.
2018-05-08	10:54:34.51	148.946	-20.030	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 36 km NE Airlie Beach Reviewed 2018-05-10.
2018-05-02	18:36:19.10	151.540	-25.217	10	1.0	Mt Perry	11 km SW Mt Perry. Reviewed 2018-05-03.
2018-04-27	18:01:08.77	151.563	-25.439	10	3.2	Gayndah	21 km N Gayndah. Reviewed 2018-06-01.
2018-04-23	04:43:54.20	148.836	-19.989	10	2.1	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 33 km NE Airlie Beach. Reviewed 2018-09-23.
2018-04-22	07:02:21.08	151.378	-25.573	10	1.9	Mundubbera	8 km E Mundubbera. Reviewed 2018-04-28.
2018-04-19	13:35:59.54	150.790	-25.643	10	2.7	Nerangy	51 km W Mundubbera. At Nerangy homestead. Heard and felt at Nerangy homestead. Reviewed 2018-09-06.
2018-04-11	05:57:33.81	147.692	-20.084	10	2.0	Bowen	Related to the 2011-04-16 05:31 ML5.3 earthquake? 59 km W Bowen. Reviewed 2018-03-24.
2018-04-10	08:57:20.63	148.721	-19.610	10	2.2	Bowen	Aftershock of 2016-08-18 04:30 event. 66 km NE Bowen. Reviewed 2018-04-11.
2018-04-05	11:27:49.73	148.841	-20.063	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 26 km NE Airlie Beach. Reviewed 2018-05-07.
2018-04-05	09:51:24.38	148.763	-19.992	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 31 km N Airlie Beach. Reviewed 2018-05-07.
2018-04-05	06:15:00.46	152.198	-24.557	10	0.9	Rosedale	30 km E Rosedale. Reviewed 2018-04-05.
2018-03-31	14:12:51.93	151.551	-25.263	10	1.7	Mt Perry	13 km SW Mt Perry. Reviewed 2018-04-05.
2018-03-29	20:41:15.87	150.362	-25.347	10	1.7	Cracow	8 km SE Cracow. Reviewed 2018-04-05.

Date (UTC)	Time (UTC)	Longitude	Latitude	Depth (km)	Magnitude (ML)	Place	Comment
2018-03-26	08:30:15.01	150.381	-25.661	10	1.3	Cracow	41 km S Cracow. Reviewed 2018-06-26.
2018-03-03	14:12:57.08	153.096	-23.068	10	2.5	Off Gladstone	206 km NE Gladstone. Reviewed 2018-03-04.
2018-02-22	06:38:36.72	144.480	-27.359	64	3.2	Yowah	69 km N Yowah. Reviewed 2018-02-22.
2018-02-21	20:30:20.42	152.872	-25.079	10	2.6	Hervey Bay	24 km N Hervey Bay. Reviewed 2018-02-22.
2018-02-18	19:21:55.83	148.775	-19.917	10	2.2	Bowen	Aftershock of 2016-08-18 04:30 event. 56 km E Bowen. Reviewed 2018-02-19.
2018-02-16	22:25:13.45	148.784	-19.971	10	1.8	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 33 km N Airlie Beach. Reviewed 2018-02-17.
2018-02-10	16:47:04.75	148.780	-20.105	10	2.6	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 19 km N Airlie Beach. Reviewed 2018-02-11.
2018-02-05	13:16:07.06	151.680	-24.756	10	1.9	Gin Gin	38 km NNW Gin Gin. Reviewed 2018-02-06.
2018-02-05	09:47:47.89	151.675	-24.781	10	1.4	Gin Gin	37 km NNW Gin Gin. Reviewed 2018-02-05. Foreshock of the 13:361.9.
2018-02-04	18:37:01.81	148.801	-20.126	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 18 km NE Airlie Beach. Reviewed 2018-09-23.
2018-02-03	03:43:10.48	148.770	-19.870	10	2.8	Bowen	Aftershock of 2016-08-18 04:30 event. 57 km NE Bowen. Reviewed 2018-02-03.
2018-01-25	17:37:02.44	148.739	-19.742	10	2.2	Bowen	Aftershock of 2016-08-18 04:30 event. 60 km NE Bowen. Reviewed 2018-01-28.
2018-01-24	02:28:16.88	153.047	-30.647	10	4.2	Nambucca Heads	4 km E Nambucca Heads. Reviewed 2018-01-24.
2018-01-15	05:28:50.86	151.551	-25.213	10	1.0	Mt Perry	10 km SW Mt Perry. Reviewed 2018-01-22.
2018-01-09	15:07:37.09	148.831	-20.074	10	1.7	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 24 km NE Airlie Beach. Reviewed 2018-01-10.
2018-01-09	10:24:25.89	148.775	-20.160	10	2.3	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 13 km NE Airlie Beach. Reviewed 2018-01-10.
2018-01-08	18:53:49.95	146.400	-16.943	10	1.8	Cairns	57 km E Cairns. Reviewed 2018-01-09.
2018-01-08	15:19:41.41	146.195	-17.147	10	1.8	Cairns	52 km SE Cairns. Reviewed 2018-01-09.

<b>Date (UTC)</b>	<b>Time (UTC)</b>	<b>Longitude</b>	<b>Latitude</b>	<b>Depth (km)</b>	<b>Magnitude (ML)</b>	<b>Place</b>	<b>Comment</b>
2018-01-08	14:43:03.03	148.715	-20.248	10	2.0	Airlie Beach	Aftershock of 2016-08-18 04:30 event. 2 km N Airlie Beach. Reviewed 2018-01-10.
2018-01-08	08:18:30.34	151.847	-25.113	10	0.8	Horse Camp	2 km SW FS03. Reviewed 2018-01-09.
2018-01-06	18:03:03.68	152.755	-24.803	10	1.3	Bundaberg	42 km ENE Bundaberg. Reviewed 2018-01-07.
2018-01-01	16:51:57.53	152.383	-18.909	10	2.8	Coral Sea	450 km NE Bowen. Reviewed 2017-01-07.

## 2018 Statistical Summary




Figure 5: Magnitude frequency count for earthquakes in the main CQSRG 2018 Catalogue.

Figure 5 provides a graphical representation of the frequency of magnitude spread for earthquake events listed in the main CQSRG Catalogue (Table 2).




Figure 6: Magnitude frequency count for earthquakes in the supplemented CQSRG 2018 Catalogue, which includes the unlocated Bowen earthquakes.

Figure 6 provides a graphical representation of the frequency of magnitude spread **for earthquake events listed in both the main CQSRG Catalogue (Table 2) and the supplementary Catalogue (Table 1)**. This shows the additional 159 unlocated earthquake events that were detected in the Whitsunday Passage, but omitted from the main CQSRG Earthquake Catalogue.





Figure 7: Yearly count of earthquakes detected by CQSRG.

Figure 7 shows the number of earthquake events detected and catalogued by CQSRG since 2004. It puts into context the extraordinary number of earthquakes detected during the past four years when compared to the numbers detected in previous years. This increase in seismic activity is entirely due to the large magnitude events and their dependent aftershocks that occurred in the Mt Perry, Rainbow Beach, and Bowen areas during 2015 and 2016.

The four past years have each exhibited a significantly greater number of earthquake events than all previous years – certainly since 2004, when CQSRG began monitoring. The 2016 ML 5.8 Bowen event is probably the second largest earthquake to have been recorded on the East coast of Mainland Australia in modern times.

## 2018 Earthquake Maps



**Figure 8: Broad view of earthquakes located in 2018 by CQSRG in Queensland.**

Figure 8 is a broad-view map of the earthquakes location in Queensland by CQSRG in 2018.

It should be noted that the earthquakes located in the South-West and North-East of the State were not directly detected by CQSRG monitoring stations.

The events in the North-East (in the Bowen area, Cairns area, and in the Coral Sea) were detected and located as part of the ongoing CQSRG study of the Bowen 2016 aftershock and reactivation sequence; using the Geoscience Australia stations BW1H and BW2S.

The two events in the South-West region were reported by Geoscience Australia and CQSRG subsequently gathered data from non-CQSRG monitoring stations to perform locational analysis.

It should not be inferred from the clustering and distributed nature of the earthquake events shown in Figure 8 that small earthquake events did not occur in other parts of Queensland. However, no

other Queensland earthquake events other than those shown in Figure 8 were reported by Geoscience Australia during the 2018 calendar year.

Geoscience Australia usually only reports on earthquakes of ML 3.5 or greater, or lesser magnitude events that have generated public interest.



## Public Seismic Network (PSN)

Since 2011-08-05 CQSRG has hosted a PSN seismograph station, known to the Australian PSN community as the Gin Gin or the Horse Camp station. Vic Dent and Mike Turnbull originally installed the station with a rudimentary setup consisting of a 3D geophone attached to a PSN A/D board, in a vacant brick shed on Mike Turnbull's property at Horse Camp, 16 km SW of Gin Gin. Mike provided a desktop computer onto which the PSN software was installed. The station regularly uploading GIF pictures of the daily seismogram traces to the Regional Seismic Users web site operated by Dale Hardy; until Dale's untimely death. That web site has now been decommissioned.

Since 2017 CQSRG has hosted a web site for the Australian Public Seismic Network at <http://cqsrq.org/psn/stations/>.

The Horse Camp station also uploads continuous data to the Regional Seismic Network (RSN), operated by the Australian Centre for Geomechanics (AGC). (Information at <https://acg.uwa.edu.au/>)

In 2013 the geophone was replaced with a Sprengnether S6000 seismometer, and the PSN A/D board was housed in a respectable electronics housing, along with custom made adaptor electronics to accommodate the sensor and GPS interface.

Since the PSN station is located only 300 m from FS03, data from the PSN station is not used in locating events detected by CQSRG, but is used to identify seismic events of interest to CQSRG.

## Appendix A – Details of FS03

Station FS03 is registered with the International Registry of Seismograph Stations maintained jointly by International Seismological Centre & World Data Centre for Seismology.

### LOCATION

Latitude -25.1068, Longitude 151.8667, Height above sea level 180 m. Approximately 16 km SW Gin Gin, Queensland, Australia.

### SITE AND SAMPLING

Sampling of ground velocity at 100 sample/sec, full scale 4194304 counts.

Ch Type	Serial	Name	Direction	Gain	Filters
1 L43D	#1482	East	90 deg true	0.00	DC 50.0
2 L43D	#1482	North	0 deg true	0.00	DC 50.0
3 L43D	#1482	Up	Positive up	0.00	DC 50.0

### DATA LOGGER

Kelunji Classic #153, GURIA V4.16A Operating System.

### TIME SYNC

Sync every day at 1400 UCT, using GPS. Wait for up to 80 seconds

Wait up to 120 seconds for a position

Auto-correct clock after sync

### TRIGGER SETTINGS

STA/LTA Channel 3, filter 1.00 to 7.50 Hz

Time const 0.20, 2.0, 20.0, 200.0 seconds

Ratios Fast 3.50, slow 1.75, squelch 5, 15 days

Length 100 to 200 secs, 80.00 sec pre-trigger, 1.10 cutoff.

## Appendix B – Details of WOOW

Station WOOW is registered with the International Registry of Seismograph Stations maintained jointly by International Seismological Centre & World Data Centre for Seismology.

### LOCATION

Latitude 25.4755, Longitude 152.0799, Height above sea level 128 m. 5 km NE of Biggenden, Queensland, Australia.

### SITE AND SAMPLING

Sampling At 100 sample/sec, full scale 4194304

Ch Type	Serial	Name	Direction	Gain	Filters
1 S6000	#10509	East	90 deg mag	1.00	DC 25.0
2 S6000	#10509	North	0 deg mag	1.00	DC 25.0
3 S6000	#10509	Up	Positive up	1.00	DC 25.0

### DATA LOGGER

Kelunji Classic #153, GURIA V4.16A Operating System.

### TIME SYNC

Sync every day at 1400 UCT, using GPS. Wait for up to 80 seconds

Wait up to 120 seconds for a position

Auto-correct clock after sync

### TRIGGER SETTINGS

TA/LTA Channel 3, filter 2.00 to 7.50 Hz

Time const 0.20, 2.0, 20.0, 200.0 seconds

Ratios Fast 2.50, slow 1.75, squelch 5, 7 days

Length 150 to 230 secs, 99.00 sec pre-trigger, 1.10 cutoff.

## Appendix C – CQSRG Method of Earthquake Location

In general, CQSRG only catalogues earthquake events that are detected by its seismic monitoring station(s). However, in the event of significant local events that, for reasons of station downtime, are not recorded by CQSRG stations, locations are conducted by obtaining data from other agencies.

The general process for earthquake event location at CQSRG is as follows.

1. Identify local earthquake events from visual inspection of CQSRG network seismograms.
2. Download extra seismograms from other agencies; typically, University of Queensland, Geoscience Australia, and the Australian National University (ANU) Australian Seismometers in Schools (AuSIS) project.
3. Send email requests to other agencies; typically, the Seismology Research Centre (SRC), and the South East Queensland Water Company (SeqWater).
4. Collect all available seismogram records and pick P and S phase arrival times using EqWave (SRC sourced software).
5. Enter the picked P and S times into EQLOCL (SRC sourced software).
6. Use the location calculated by EQLOCL.

In the not so rare cases where the only record available is that from FS03, an attempt is made to locate the event using first motion polarity and near field trigonometry. This can only be done when the first motions are sufficiently impulsive to give an unambiguous indication of the arrival azimuth. In cases where only two records are available (invariably FS03 and EIDS), and the S-P derived radial distance circles meet, but do not over extend, the touch point is used as a seed to the EQLOCL algorithm.

In cases where only two records are available (invariably FS03 and EIDS), and the S-P derived radial distance circles over extend, but the first motions are sufficiently impulsive to derive an unambiguous azimuth, the radial touch point indicated by the azimuth direction is used as a seed to the EQLOCL algorithm.

In cases where only two records are available (invariably FS03 and EIDS), and the S-P derived radial distance circles over extend, but the first motions are insufficiently impulsive to derive an unambiguous azimuth, the locations of both the radial touch points are used as seeds to the EQLOCL algorithm, and the resulting ambiguous locations are noted in the catalogue entry comments.

In cases where the only information that can be gleaned is the radial distance from FS03, that distance may be noted in the catalogue listing comments.

## Appendix D – CQSRG Method of Magnitude Quantification from FS03 Records

### Calibration of FS03 Seismometer for Earthquake Magnitude Determination.

Mike Turnbull, 7 November, 2012.

#### Introduction

FS03 is the designation of a seismic monitoring station operated by the Central Queensland Seismology Research Group (CQSRG). It is located about 16 km south-west of Gin Gin.

When the FS03 station was first installed it had a Sprengnether S6000 seismometer attached to a data logger manufactured by the Seismology Research Centre (SRC). The characteristics of this sensor and the amplification factors of the data logger section of the seismograph were used as input to the SRC software used to locate and quantify earthquakes recorded on the seismograph. When the S6000 sensor failed it was replaced with a Mark Products L43D seismometer sensor. By comparison of the calibration waveform amplitudes of the S6000 against the L43D, a correction factor of 1.7 was calculated and used to adjust the amplitude value input to the SRC software to determine earthquake magnitudes using the new sensor – and this provided a temporary solution.

In order for the SRC software to be able to calculate an earthquake magnitude, it first must be able to calculate the earthquake's epicentral location. This can only be done if seismographic records from at least three different stations are available. In situations where only one or two records are available the software cannot locate the epicentre. Consequently, in cases where an earthquake cannot be located, determination of its magnitude using EQLOCL has always been problematic.

This appendix describes a method of extracting parametric information from past earthquake magnitudes, located with the SRC software using FS03 seismograms, that can be used in a suitable mathematical formula to determine the magnitude of other earthquakes recorded on the FS03 seismograph, using information from the single station data. This allows the magnitude determination to be done independent of the SRC software.

#### Background Information

The Richter local earthquake magnitude ( $M$ ) is calculated using the formula given in Eq. 1.

$$M = \log_{10}A - \log_{10}A_0 \text{ (Eq. 1)}$$

Where:

$A$  is the maximum amplitude of the seismic record of the earthquake, and

$A_0$  is the maximum amplitude that would be produced on the same sensor by an earthquake of magnitude zero, occurring at the same location as the earthquake under consideration.

The value of  $\log_{10}A_0$  is dependent only on the epicentral distance of the earthquake from the sensor, and the response characteristics of the sensor itself. It is assumed that the relationship is as given in Eq. 2 (**NOTE: This assumed relationship has yet to be confirmed as being valid**).

$$\log_{10}A_0 = a\delta + b \quad \text{(Eq. 2)}$$

Where:

$\delta$  is the epicentral distance, and  
 a and b are parameters yet to be determined, characteristic of the sensor.

### Method

It is clear that Eq.2 is linear. Therefore the sensor parameters a and b can be determined from the slope and intercept, respectively, of the graph of  $\log_{10}A_0$  plotted against  $\delta$ , providing that sufficient data is available

The epicentral distance  $\delta$  can be expressed in any value that provides a valid determination of the distance from the sensor to the epicentre. This could be (for example):

- the difference in arrival times of the P and S waves (in seconds for example); or,
- the surface distance from sensor to epicentre (in km for example); or,
- the Earth centric angle of arc from sensor to epicentre (in degrees for example).

The values for  $\log_{10}A_0$  can be calculated from past earthquake events, the magnitudes of which have been determined with the SRC software using FS03 seismograms.

Transformation of Eq.1 gives Eq. 3.

$$\log_{10}A_0 = \log_{10}A - M \text{ (Eq.3)}$$

Table 3 presents the calculations of  $\log_{10}A_0$  based on nine past events that were quantified with the SRC software, showing the S-P time differences used to measure epicentral distances.

**Table 3: Determination of  $\log_{10}A_0$  from past events recorded on the FS03 seismograph.**

Earthquake Date	Measured P arrival in relative seconds	Measured S arrival in relative seconds	S-P time (s)	Measured Amplitude A	Magnitude estimated using ES&S algorithm M	Calculated $\log_{10}(A_0)$
2012-09-19 06:14	11.54	14.97	3.43	1900	1.6	1.6787536
2012-05-20 17:58	42.23	45.79	3.56	1103	1.5	1.5425755
2012-09-22 23:59	38.31	41.91	3.6	243	1.0	1.3856063
2012-04-10 01:51	37.54	42.57	5.03	473.2	1.4	1.2750447
2012-09-25 03:06	10.56	22.7	12.14	456	1.9	0.7589648
2012-08-19 22:37	29.38	41.82	12.44	215	1.5	0.8324385
2012-09-03 15:04	10.86	26.84	15.98	1828	2.8	0.4619762
2012-09-23 16:29	36.21	53.77	17.56	3620	3.2	0.3587086
2012-01-05 14:05	9.96	56.18	46.22	1352	4.3	-1.1690233

Figure 9 shows the graph of  $\log_{10}A_0$  plotted against the associated S-P time difference (extracted from Table 3).




Figure 9:  $\log_{10}A_0$  Vs S-P.

Figure 9 also displays the line of best fit, calculated using linear regression of the plotting data, along with the slope, intercept, and correlation coefficient ( $R^2$ ). **The  $R^2$  value of 0.97 confirms that the assumed linear relationship is valid.**

By substituting the slope and intercept values into Eq.1 and Eq.2 we arrive at the formula for FS03 magnitudes given in Eq.4.

$$M_{FS03} = \log_{10}A - (-0.064(S - P) + 1.64) \quad (\text{Eq.4})$$

Where:

$M_{FS03}$  is the Richter magnitude determined from an FS03 seismogram record;

A is the maximum amplitude of the unfiltered FS03 seismogram record;

S is the arrival time of the S wave in seconds, and;

P is the arrival time of the P wave in seconds.

### Important Note Concerning Accuracy and Precision

Table 3, Figure 9, and Equation 4, show a shortened calculation using only 9 historical events, to demonstrate the method. A consequence of using so few input values is that the resulting error ranges will suffer. Consequently, in order to reduce the standard errors in magnitude calculations based on this method, and extend the accuracy to at least one decimal point, many more input data are required.

The calculations used to determine the actual  $\log_{10}A_0$  values for FS03, used in quantifying earthquake magnitudes, used 34 historical events. This resulted in parameter **a** and **b** values for Equation 2, as shown in Table 4.


**Table 4: Equation 2, a and b Parameter values and Standard Errors.**

	<b>a</b>	<b>b</b>
<b>Estimation</b>	-0.088	1.81
<b>Standard Error</b>	±0.004	±0.05
<b>Correlation</b>	0.94	

This implies that magnitudes determined using this method will be accurate to at least one decimal place. The a and b values shown in Table 4 are those used at CQSRG to calculate local magnitudes of events recorded by station FS03.

### Example Usage

Figure 10 shows the seismogram of an earthquake that was recorded on station FS03 on 26 October 2012.



**Figure 10: FS03 record of an earthquake.**

From Figure 10 we can obtain the maximum amplitude ( $A = 460$ ), the P wave arrival time ( $P = 25.69$  s) and the S wave arrival time ( $S = 27.90$  s); from which the time difference ( $S - P = 2.21$  s) can be determined.

Inserting these values into Eq.4 we calculate a Richter magnitude of 1.2 (rounded to one decimal place).

Table 5 shows the results of some other similar calculations, for different earthquakes.



Table 5: Calculations of FS03 Richter magnitudes for some earthquakes.

Earthquake Date	Measured P arrival in relative seconds	Measured S arrival in relative seconds	S-P time (s)	Measured Amplitude A	Calculated $M_{FS03}$ Magnitude
2012-09-28 16:38	10.56	22.7	12.14	304	1.6
2012-10-03 17:29	25.09	27.69	2.6	259	0.9
2012-10-18 14:48	23.43	25.91	2.48	911	1.5
2012-10-26 04:47	25.69	27.9	2.21	460	1.2

### Student Resources

Figure 11, Figure 12, Figure 13, and Figure 14 are images of earthquake seismograms recorded by FS03. They are included here for the reader to use as practice on the CQSRG magnitude determination method. They can also be used as a resource for High School science teachers who may want to use the formulae presented here as real-world examples of applied mathematics.




Figure 11: Earthquake recorded on FS03 on 28 September 2012.




Figure 12: Earthquake recorded on FS03 on 3 October 2012.




Figure 13: Earthquake recorded on FS03 on 18 October 2012.




Figure 14: Earthquake recorded on FS03 on 26 October 2012.

## Appendix E - CQSRG Method of Magnitude Quantification from EIDS Records

### Relative Calibration of EIDS Seismometer for Earthquake Magnitude Determination Based on FS03 Past events.

Mike Turnbull, 17 Feb, 2015.

#### Introduction

FS03 is the designation of a seismic monitoring station operated by the Central Queensland Seismology Research Group (CQSRG). It is located about 24 km south-west of Gin Gin.

EIDS is the Geoscience Australia station located near Eidsvold. The characteristics of the EIDS sensor and associated equipment are unknown (to the author); however, it would be useful to be able to estimate event magnitudes using records from EIDS.

This paper describes a method of extracting parametric information from past earthquakes recorded by both FS03 and EIDS, and quantified using the FS03 seismograms or some other reliable method, that can be used in a suitable mathematical formula to determine the magnitude of earthquakes recorded on the EIDS seismograph.

#### Background Information

The Richter local earthquake magnitude ( $M$ ) is calculated using the formula given in Eq. 1.

$$M = \log_{10}A - \log_{10}A_0 \text{ (Eq. 1)}$$

Where:

$A$  is the maximum amplitude of the seismic record of the earthquake on a given sensor, and  $A_0$  is the maximum amplitude that would be produced on the same sensor by an earthquake of magnitude zero, occurring at the same location as the earthquake under consideration.

The value of  $\log_{10}A_0$  is dependent only on the epicentral distance of the earthquake from the sensor, and the response characteristics of the sensor itself. It is assumed that the relationship is linear as given in Eq. 2 (**NOTE: This assumed relationship has yet to be confirmed as being reasonable**).

$$\log_{10}A_0 = a\delta + b \quad \text{(Eq. 2)}$$

Where:

$\delta$  is the epicentral distance from the sensor under consideration, and  $a$  and  $b$  are parameters yet to be determined, characteristic of the sensor under consideration.

#### Method

Eq.2 is linear, therefore the sensor parameters  $a$  and  $b$  can be determined from the slope and intercept, respectively, of the graph of  $\log_{10}A_0$  plotted against  $\delta$ , using linear regression, providing that sufficient data is available for the sensor being considered.

The epicentral distance  $\delta$  can be expressed in any value that provides a valid determination of the distance from the sensor to the epicentre. This could be (for example):

- the difference in arrival times of the P and S waves (in seconds for example); or,
- the surface distance from sensor to epicentre (in km for example); or,
- the Earth centric angle of arc from sensor to epicentre (in degrees for example).

The values for  $\log_{10}A_0$ , for the sensor under consideration, can be calculated from the amplitudes and S-P times of records of past earthquake events, the magnitudes of which events have been determined by some other reliable method – in this case, from magnitudes determined from FS03 records, or as published by Geoscience Australia.

Transformation of Eq.1 gives Eq. 3.

$$\log_{10}A_0 = \log_{10}A - M \text{ (Eq.3)}$$

Table 6 presents the calculations of  $\log_{10}A_0$  values for EIDS based on past events that were quantified with FS03 seismograms, showing the S-P time differences used to measure epicentral distances from the EIDS sensor. The EIDS seismograms were all similarly conditioned using a 2 Hz to 10 Hz band-pass filter.

**Table 6: Determination of  $\log_{10}A_0$  from past events recorded on the EIDS seismograph.**

Earthquake Date	Measured EIDS P arrival	Measured EIDS S arrival	EIDS S-P	Measured EIDS Amplitude A	Magnitude Estimated using FS03 M	Calculated EIDS $\log_{10}(A_0)$
14/06/2014 14:19	17.82	25.48	7.66	99198	3	1.996503
26/06/2014 11:02	17.63	45.34	27.71	678	2.4	0.43123
22/08/2014 08:34	37.56	43.62	6.06	6649	1.9	1.922756
22/08/2014 08:35	27.14	33.69	6.55	92100	2.7	2.26426
22/08/2014 08:38	21.63	28.42	6.79	137217	2.8	2.337408
03/01/2013 19:11	55.16	65.87	10.71	1422	1.6	1.5529
07/01/2013 18:41	60.16	71.28	11.12	822	1.3	1.614872
14/02/2013 23:03	15.7	34.76	19.06	3992	2.1	1.501191
05/01/2012 14:05	75.3	128.37	53.07	4691	4.3	-0.62873
10/04/2012 01:51	43.44	52.12	8.68	2440	1.4	1.98739
20/05/2012 17:58	50.12	59.56	9.44	3879	1.5	2.08872
19/08/2012 22:37	16.48	19.93	3.45	14340	1.5	2.656549
03/09/2012 15:03	15.13	34.44	19.31	14467	2.8	1.360378
19/09/2012 06:14	17.89	25.65	7.76	5905	1.6	2.17122
22/09/2012 23:59	40.45	46.17	5.72	1761	1	2.245759
23/09/2012 16:29	32.05	46.23	14.18	65547	3.2	1.616553
25/09/2012 03:05	22.88	44.51	21.63	2944	1.9	1.568938
03/12/2012 07:41	44.89	51.7	2447	9544	1.4	2.57973
04/12/2012 20:17	55.01	61.41	6.4	570212	3.2	2.556036

12/12/2012 10:36	53.69	60.33	6.64	20513	2.2	2.112029
------------------	-------	-------	------	-------	-----	----------

Figure 15 shows the graph of  $\log_{10}A_0$  plotted against the associated S-P time difference (extracted from Table 6: Determination of  $\log_{10}A_0$  from past events recorded on the EIDS seismograph. Figure 15 also displays the line of best fit, calculated using linear regression of the plotting data, along with the slope, intercept, and correlation coefficient ( $R^2$ ). **The  $R^2$  value of 0.91 confirms that the assumed linear relationship is reasonably valid.**




Figure 15: 23:  $\log_{10}A_0$  Vs S-P

By substituting the slope and intercept values into Eq.1 and Eq.2 we arrive at the formula for EIDS magnitudes given in Eq.4.

$$M_{EIDS} = \log_{10}A - (-0.064(S - P) + 2.63) \quad (\text{Eq.4})$$

Where:

- $M_{EIDS}$  is the Richter magnitude determined from an EIDS seismogram record;
- A is the maximum amplitude of the EIDS seismogram record;
- S is the arrival time of the S wave in seconds, and;
- P is the arrival time of the P wave in seconds.

### Example Usage

Figure 16 shows the seismogram of an earthquake that was recorded on station EIDS on 15 February 2015.




Figure 16: EIDS record of an earthquake.

From Figure 16 we can obtain the maximum amplitude ( $A = 13793261$ ), the P wave arrival time ( $P = 17.03$  s) and the S wave arrival time ( $S = 22.93$  s); from which the time difference ( $S - P = 22.93$  s) can be determined.

Inserting these values into Eq.4 we calculate a Richter magnitude of 4.9 (rounded to one decimal place).

Table 7 shows the results of some other similar calculations, for different earthquakes, along with the GA published magnitudes for the same events.

Table 7: Calculations of EIDS Richter magnitudes for some earthquakes.

Earthquake Date	Measured P arrival	Measured S arrival	S-P	EIDS Amplitude A	Calculated $M_{EIDS}$ Magnitude	GA Published Magnitude
15/02/2015 15:57	17.03	22.93	5.9	13793261	4.9	5.1
15/02/2015 15:58	12.58	18.56	5.98	869195	3.7	
15/02/2015 16:40	43.7	49.25	5.55	278525	3.2	3.2
15/02/2015 17:37	13.18	19.15	5.97	907859	3.7	3.4
15/02/2015 18:06	14.56	20.54	5.98	125151	2.9	2.5
16/02/2015 05:56	58.18	64.14	5.96	1703875	4.0	4.0

## Appendix F - Magnitude Calibration of BW1H for the Bowen 2016 Earthquake Sequence.

### Introduction

In August 2016 a magnitude 5.8 earthquake occurred in the Whitsunday Passage east of Bowen and North of Airlie Beach. This was followed by many aftershocks over the next few months. The nearest seismic monitoring stations were the Urban Monitoring (UM) network stations in Bowen. There are two stations: one on soft basement (BW2S); and one on hard basement (BW1H).

The BW1H station provided relatively clean records of the main and aftershocks.

The locations of the earthquakes in the sequence formed a relatively tight group around the main event. Figure 17 shows the grouping pattern.




Figure 17: The grouping of the 2016 Bowen earthquake sequence.

The main event was located 64 km NNE of Bowen, with the other earthquakes ranging from 40 km to 100 km radial distance from Bowen.

The relative proximity of all earthquakes to Bowen meant that the radial attenuation of the seismic energy over that distance would vary only slightly; and it was assumed that any such slight attenuation would not adversely affect any logarithmic magnitude values determined within the 40 to 100 km range. It was therefore decided to calibrate the BW1H records against the Geoscience Australia (GA) published magnitudes, so that those records could be used to determine the magnitudes of those smaller events not analysed by GA.



The method of calibration was as follows.

### Method of Calibration

Thirty three earthquakes with published GA magnitudes in the range from ML 2.1 to ML 5.8 were selected (See Table 8).

**Table 8: List of earthquake magnitudes published by GA.**

Date	Time	P	S	S-P	Amplitude 2-10 Hz	Published GA Mag	Calculated BW1H ML
2016-08-20	01:17	4654.46	4662.07	7.61	669	2.2	2.2
2016-08-19	17:03	61422.39	61430.16	7.77	697	2.1	2.2
2016-08-18	15:47	56863.64	56871.33	7.69	753	2.3	2.2
2016-08-19	22:41	81707.73	81715.41	7.68	1205	2.5	2.4
2016-08-18	17:17	62271.91	62279.60	7.69	1324	2.4	2.4
2016-08-19	11:15	40552.57	40560.20	7.63	1574	2.2	2.4
2016-08-20	20:11	72668.06	72675.76	7.70	1794	2.3	2.5
2016-08-18	15:05	54340.74	54348.38	7.64	1810	2.6	2.5
2016-08-20	00:53	3224.43	3232.12	7.69	1857	2.1	2.5
2016-08-18	13:57	50252.43	50260.39	7.96	2159	2.3	2.5
2016-08-19	23:56	86213.38	86220.91	7.53	2349	2.4	2.6
2016-08-18	16:23	58987.97	58995.55	7.58	2387	2.8	2.6
2016-08-18	05:54	21249.73	21257.29	7.56	2430	2.7	2.6
2016-08-18	10:06	36414.70	36422.36	7.66	2653	2.7	2.6
2016-08-19	20:40	74439.78	74447.37	7.59	2715	2.4	2.6
2016-08-18	15:52	57131.48	57138.86	7.38	2928	2.5	2.7
2016-08-18	07:35	27311.40	27319.08	7.68	2940	2.6	2.7
2016-08-18	09:23	33823.41	33830.84	7.43	2965	2.8	2.7
2016-08-18	05:57	21467.21	21474.69	7.48	3240	2.6	2.7
2016-08-18	05:36	20196.43	20203.14	6.71	3561	3.3	2.7
2016-08-20	07:10	25849.39	25857.21	7.82	3832	2.8	2.8
2016-08-18	21:38	77887.80	77895.31	7.51	4140	3.1	2.8
2016-08-20	16:46	60410.48	60418.09	7.61	9503	3.2	3.1
2016-08-18	05:09	18579.10	18586.66	7.56	9553	2.9	3.1
2016-08-18	14:03	50598.61	50606.19	7.58	9780	3.4	3.1
2016-08-18	04:36:53.28	16622.15	16629.53	7.38	10228	3.9	3.1
2016-08-18	05:23	19445.18	19452.59	7.41	22102	3.6	3.5
2016-08-18	09:30	34261.85	34269.63	7.78	22753	3.5	3.5
2016-08-18	08:56	32219.57	32227.10	7.53	40768	3.4	3.8
2016-08-18	04:39:52.05	16740.36	16747.92	7.56	42854	3.8	3.8
2016-08-18	05:30	19847.18	19854.69	7.51	52254	4.0	3.9
2016-08-18	18:27	66467.63	66474.96	7.33	128115	4.1	4.4
2016-08-18	04:30:08.43	16219.32	16227.00	7.68	973331	5.8	5.8

The amplitudes stated in Table 8 are those of the associated earthquake's S phase maximum, with a bandpass filter of 2 Hz to 10 Hz applied to the time series data.

The filtered amplitudes were then plotted against the published GA magnitudes as shown in Figure 18.




Figure 18: Filtered Amplitudes Vs Published Magnitudes.

A power curve was then fitted to the plotted data to determine the following relation.

$$M_{BW1H} = 0.9019A_{BW1H}^{0.1353}$$

This relation has a correlation coefficient of 87%.

Although the use of the power relation parameters to four decimal places is not warranted (two decimal places would probably suffice), the full precision was maintained when determining the magnitudes of 2016 Bowen aftershocks not published by GA.

### Verification of Calibration

The right hand column of Table 8 lists the  $M_{BW1H}$  values calculated using the power relation. This shows that the calculated magnitudes are typically accurate to one decimal place  $\pm 0.1$ , in the range from ML 2.0 up to ML 5.8, with occasional inaccuracies up to  $\pm 0.8$ .

### Calibration Saturation

Inspection of Figure 18 indicates that the power relation is flattening out at the bottom end to become asymptotic between ML 2.0 and ML 1.5.

Therefore  $M_{BW1H}$  values in the range ML 1.5 to ML 2.0 will be overestimated, and values below that range should be ignored.

## Appendix G - Method Used to Identify Bowen August 2016 ML 5.8 Aftershocks

As state in the CQSRG web page at <http://cqsrq.org/>, CQSRG's primary research aim is to monitor for, and catalogue, earthquakes in Eastern Central Queensland; in the region bounded (approximately) north to Mackay, South to the Sunshine Coast, west to Roma, and out to sea some hundreds of kilometres. The main reason for generally restricting research to that broad region is that it encompasses the earthquake detection capability of the main recording station operated by CQSRG – that is, the FS03 seismic monitoring station just west of Gin Gin, in the Bundaberg Regional area. When the Bowen August 2016 ML 5.8 event occurred it triggered the FS03 recorder, as did several of the aftershocks that occurred in the following weeks.

As the aftershock sequence progressed, the average magnitude of the events reduced. Consequently, after a short time, the aftershocks were no longer triggering on the CQSRG network. However, experience gained during analysis of the earlier, larger magnitude aftershocks indicated that the lesser magnitude aftershocks, of about ML 1.3 and above, could be unambiguously identified by visual inspection of the BW1H and BW2s station seismograms with a very high degree of reliability.

The key diagnostic features used to visually identify the aftershock recordings were:

- The consistent S-P times being confined to a precise spread within three standard deviations of the sequence average.
- The characteristic shape of the wave form (the wave form *signature*).
- Suitable and consistent choice of amplitude and time-scale gain settings on the seismogram viewer being used to visually inspect the seismogram records.

### Analysis of S-P Times

In the first two days following the main earthquake Geoscience Australia published location solutions for 34 events (including the main event and aftershocks). The closest station to the events was BW1H, the Bowen hard site Queensland UMP station. Using the S and P arrival times picked from the BW1H records of the 34 published events the S-P times were obtained, averaged, and the sample standard deviation was derived. The results of this analysis are provided in Table 9.

Table 9: Statistical Analysis of Main event and aftershock S-P Times.

S-P Time	Statistic Description
7.6	Average S-P time for 34 events recorded on BW1H and verified by Geoscience Australia.
0.2	Sample standard deviation of the aftershock S-P times.
8.2	Upper S-P time expected for valid aftershocks.
6.9	Lower S-P time expected for valid aftershocks.

Based on the statistics listed in Table 9, individual events with S-P times greater than 8.2 s were treated with suspicion as being non-aftershock events. Extra analysis and inspection of the vast

majority of these suspect events proved them to be exclusively extraction blasts from the Collinsville and Sonoma Coal Mines, 70 km south west of Bowen.

Many of the prospective “events” with S-P times less than 6.9 s, especially those with dramatically short S-P times, turned out on further inspection and analysis to be local social noise – as shown by not being present on the BW2S records. However, valid aftershock events were recorded with S-P times as low as 5.59 s.




Figure 19: BW1H and BW2S Recordings of the Main ML 5.8 event.

### The characteristic shape of the wave form

Figure 19 is the seismographic record of the main ML 5.8 earthquake event, as recorded on the BW1H and BW2S stations’ vertical sensors, and as viewed on the Seismology Research Centre’s (SRC) Waves V1.2 Seismic Waveform Analysis Software. The visual characteristics to note are:

- The S-P time, as previously mentioned.
- The impulsive nature of both the S and P arrivals.
- The rapid decay of the P train resulting in a single maximum amplitude P peak, and clean distinction of the subsequent S arrival.
- The rapid decay of the S train resulting in a single vertical line maximum S amplitude feature.

Observation of numerous verified aftershock records showed that these visual characteristics were invariably present in all valid aftershock records – even those of magnitudes less than ML 1.5; as can be seen in Figure 20, a BW1H/BW2S recording of an ML 1.3 aftershock.



Figure 20: Record of an ML 1.3 Aftershock on BW1H and BW2S.

Despite the signature of the P and S arrivals on both BW1H and BW2S stations being only just proud of the ambient noise, they can be readily distinguished on both stations (indicating that it is not a local noise event), and the S-P times for both stations are consistent with the expectation statistics. The magnitude of ML 1.3 was observed to be the lowest magnitude aftershock recording that could be reliably discriminated from the ambient background noise. However, it is considered that even the ML 1.3 events would only have been detected under ideal conditions, and that many valid aftershock events of this magnitude would have not been detected by this observational method. While Figure 19 and Figure 20 represent the two extremes of the magnitude range, the following Figure 21 and Figure 22 provide examples of mid-range aftershocks of ML 1.8, ML 2.2 and ML 2.6.

All three of the events in Figure 21 and Figure 22 have been located using data from multiple stations and verified as being valid aftershocks.




Figure 21: Example of an ML 1.8 aftershock




Figure 22: Example of ML 2.2 and ML 2.6 aftershocks.